

SOME REMARKS ON THE NAVIER-STOKES EQUATIONS WITH
REGULARITY IN ONE DIRECTION

ZUJIN ZHANG, WEIJUN YUAN, Ganzhou, YONG ZHOU, Zhuhai

Received September 29, 2018. Published online April 11, 2019.

Abstract. We review the developments of the regularity criteria for the Navier-Stokes equations, and make some further improvements.

Keywords: regularity criteria; Navier-Stokes equations

MSC 2010: 35B65, 35Q30, 76D03

1. INTRODUCTION

The homogeneous incompressible fluid flow is governed by the following Navier-Stokes equations (NSE):

$$(1.1) \quad \begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} - \Delta \mathbf{u} + \nabla \pi = \mathbf{0}, \\ \nabla \cdot \mathbf{u} = 0, \\ \mathbf{u}(0) = \mathbf{u}_0, \end{cases}$$

where $\mathbf{u} = (u_1, u_2, u_3)$ is the fluid velocity field, π is a scalar pressure, \mathbf{u}_0 is the prescribed initial velocity field satisfying the compatibility condition $\nabla \cdot \mathbf{u}_0 = 0$, and

$$\partial_t \mathbf{u} = \frac{\partial \mathbf{u}}{\partial t}, \quad \partial_i = \frac{\partial}{\partial x_i}, \quad (\mathbf{u} \cdot \nabla) = \sum_{i=1}^3 u_i \partial_i.$$

The global existence of a weak solution to the evolutionary NSE (1.1) was established by Leray [12] and Hopf [8] long time ago. However, the issue of its regularity

The research has been supported by the National Natural Science Foundation of China (grant no. 11761009).

and uniqueness remains open up to now. Pioneered by Serrin [19], we began studying the regularity criterion for the NSE (1.1); that is, finding a sufficient condition for the smoothness of the solution. The classical Prodi-Serrin conditions (see [6], [17], [19]) says that if

$$(1.2) \quad \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 1, \quad 3 \leq q \leq \infty,$$

then the solution is regular on $(0, T)$.

This was generalized by Beirão da Veiga [1] by considering the velocity gradient or vorticity,

$$(1.3) \quad \nabla \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 2, \quad \frac{3}{2} \leq q \leq \infty.$$

Notice that the case $q \in [\frac{3}{2}, 3)$ follows directly from (1.2) and the Sobolev inequality.

In view of the divergence-free condition $\nabla \cdot \mathbf{u} = 0$, it is natural to ask whether or not we can reduce (1.2) and (1.3) to its partial components. One way is to consider regularity criteria involving only one velocity component, which were done in [3], [9], [10], [13], [25], [27], and the references therein. Another way is to study the possible components reduction of $\nabla \mathbf{u}$ to ∇u_3 , see for instance [4], [5], [7], [10], [16], [20], [22], [24], [26], [27]; or to $\partial_3 \mathbf{u}$, see for example [2], [11], [15], [14], [23]. In [14], Penel and Pokorný showed that if

$$(1.4) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = \frac{3}{2}, \quad 2 \leq q \leq \infty,$$

then the solution is smooth. This was improved by Kukavica-Ziane [11] to be

$$(1.5) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 2, \quad \frac{9}{4} \leq q \leq 3.$$

Later on, Cao [2] employed multiplicative Sobolev inequalities

$$(1.6) \quad 1 \leq q < \infty \Rightarrow \|f\|_{L^{3q}} \leq C \|\partial_1 f\|_{L^2}^{1/3} \|\partial_2 f\|_{L^2}^{1/3} \|\partial_3 f\|_{L^q}^{1/3}$$

and

$$(1.7) \quad 1 \leq q < \infty \Rightarrow \|f\|_{L^{5q}} \leq C \|\partial_1(f^2)\|_{L^2}^{1/5} \|\partial_2(f^2)\|_{L^2}^{1/5} \|\partial_3 f\|_{L^q}^{1/5}$$

to get the following extended regularity condition:

$$(1.8) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 2, \quad \frac{27}{16} \leq q \leq \frac{5}{2}.$$

In a recent paper, Zhang [23] generalized (1.7), and improved (1.5) and (1.8), simultaneously. Precisely, he showed that the condition

$$(1.9) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = 2, \quad 1.56207 \approx \frac{3\sqrt{37}}{4} - 3 \leq q \leq 3$$

could ensure the regularity of the solution.

Combining the progress listed above, we see that the state of the regularity criterion for the Navier-Stokes equations is the following:

$$(1.10) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = \begin{cases} 2, & \frac{3\sqrt{37}}{4} - 3 \leq q \leq 3, \\ \frac{3}{2}, & 3 < q \leq \infty. \end{cases}$$

Consequently, two questions appeared naturally. One is whether we could establish regularity conditions via $\partial_3 \mathbf{u}$ for the space integrability index

$$q \in \left(\frac{3}{2}, \frac{3\sqrt{37}}{4} - 3 \right).$$

Second is to improve the scaling dimension $\frac{3}{2}$ for $q \in [3, \infty)$.

In this paper, we shall prove the following two regularity criteria:

$$(1.11) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = \frac{8}{5} + \frac{3}{5q}, \quad 4 \leq q \leq \infty,$$

and

$$(1.12) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = \frac{14}{11} + \frac{9}{11q}, \quad \frac{5}{2} \leq q \leq \infty.$$

Whence, the up-to-date status of this topic is the following smoothness condition:

$$(1.13) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = \begin{cases} 2, & \frac{3\sqrt{37}}{4} - 3 \leq q \leq 3, \\ \frac{14}{11} + \frac{9}{11q}, & 3 < q < \frac{18}{5}, \\ \frac{3}{2}, & \frac{18}{5} \leq q < 4, \\ \frac{8}{5} + \frac{3}{5q}, & 4 \leq q \leq \infty. \end{cases}$$

Before stating the main result, let us recall the weak solution of (1.1) in the sense of Leray and Hopf, see for instance [18], Definitions 3.3 and 4.9.

Definition 1.1. Let $\mathbf{u}_0 \in L^2(\mathbb{R}^3)$ with $\nabla \cdot \mathbf{u}_0 = 0$, $T > 0$. A measurable \mathbb{R}^3 -valued function \mathbf{u} defined on $[0, T] \times \mathbb{R}^3$ is said to be a weak solution to (1.1) if

- (1) $\mathbf{u} \in L^\infty(0, T; L^2(\mathbb{R}^3)) \cap L^2(0, T; H^1(\mathbb{R}^3))$;
(2) (1.1)₁ and (1.1)₂ hold in the sense of distributions, i.e.

$$\int_0^t \int_{\mathbb{R}^3} \mathbf{u} \cdot [\partial_t \varphi + (\mathbf{u} \cdot \nabla) \varphi] dx ds + \int_{\mathbb{R}^3} \mathbf{u}_0 \cdot \varphi(0) dx = \int_0^t \int_{\mathbb{R}^3} \nabla \mathbf{u} : \nabla \varphi dx dt$$

for each $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^3)$ with $\nabla \cdot \varphi = 0$, where $A : B = \sum_{i,j=1}^3 a_{ij} b_{ij}$ for 3×3 matrices $A = (a_{ij})$, $B = (b_{ij})$, and

$$\int_0^T \int_{\mathbb{R}^3} \mathbf{u} \cdot \nabla \psi dx dt = 0$$

for each $\psi \in C_c^\infty(\mathbb{R}^3 \times [0, T])$;

- (3) the strong energy inequality, that is,

$$(1.14) \quad \|\mathbf{u}(t)\|_{L^2}^2 + 2 \int_s^t \|\nabla \mathbf{u}(\tau)\|_{L^2}^2 d\tau \leq \|\mathbf{u}(s)\|_{L^2}^2 \quad \forall s < t < T,$$

holds for $s = 0$ and almost all times $s \in (0, T)$.

Now, our main result reads:

Theorem 1.2. Let $\mathbf{u}_0 \in L^2(\mathbb{R}^3)$ with $\nabla \cdot \mathbf{u}_0 = 0$, $T > 0$. Assume that \mathbf{u} is a weak solution to (1.1) on $[0, T]$ with initial data \mathbf{u}_0 . If

$$(1.15) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = \frac{8}{5} + \frac{3}{5q}, \quad 4 \leq q \leq \infty,$$

or

$$(1.16) \quad \partial_3 \mathbf{u} \in L^p(0, T; L^q(\mathbb{R}^3)), \quad \frac{2}{p} + \frac{3}{q} = \frac{14}{11} + \frac{9}{11q}, \quad \frac{5}{2} \leq q \leq \infty,$$

then the solution \mathbf{u} is smooth on $(0, T] \times \mathbb{R}^3$.

The proof of Theorem 1.2 will be provided in Section 2. Before doing that, let us recall an interesting combinatoric regularity condition by Penel and Pokorný [14], Theorem 1 (a):

$$(1.17) \quad \begin{aligned} u_3 &\in L^{2s/(s-3)}(0, T; L^s(\mathbb{R}^3)), & 3 < s \leq \infty; \\ \partial_3 u_1, \partial_3 u_2 &\in L^{2q/(2q-3)}(0, T; L^q(\mathbb{R}^3)), & \frac{3}{2} < q \leq \infty. \end{aligned}$$

Also, two inequalities concerning the pressure are recalled (see [14], Lemma 1):

$$(1.18) \quad \|\pi\|_{L^r} \leq C \|\mathbf{u}\|_{L^{2r}}^2, \quad \|\partial_3 \pi\|_{L^r} \leq C \sum_{i,j=1}^3 \|\partial_3(u_i u_j)\|_{L^r}, \quad 1 < r < \infty.$$

2. PROOF OF THEOREM 1.2

In this section, we shall prove Theorem 1.2.

For any $\varepsilon \in (0, T)$, since $\mathbf{u} \in L^2(0, T; H^1(\mathbb{R}^3))$ and (1.14) holds for almost all times $s \in (0, T)$, we may find a $\delta \in (0, \varepsilon)$ such that

$$\mathbf{u}(\delta) = \mathbf{u}(\cdot, \delta) \in L^2(\mathbb{R}^3), \quad \nabla \mathbf{u}(\delta) \in L^2(\mathbb{R}^3) \Rightarrow \mathbf{u}(\delta) \in L^3(\mathbb{R}^3),$$

as well as

$$\|\mathbf{u}(t)\|_{L^2}^2 + 2 \int_{\delta}^t \|\nabla \mathbf{u}(\tau)\|_{L^2}^2 d\tau \leq \|\mathbf{u}(\delta)\|_{L^2}^2 \quad \forall \delta < t < T.$$

Take this $\mathbf{u}(\delta)$ as initial data, there exists an $\tilde{\mathbf{u}} \in C([\delta, \Gamma^*), H^1(\mathbb{R}^3)) \cap L^2(0, \Gamma^*; H^2(\mathbb{R}^3))$, where $[\delta, \Gamma^*)$ is the life span of the unique strong solution, see [21]. Moreover, $\tilde{\mathbf{u}} \in C^\infty(\mathbb{R}^3 \times (\delta, \Gamma^*))$. According to the uniqueness result, $\tilde{\mathbf{u}} = \mathbf{u}$ on $[\delta, \Gamma^*)$. If $\Gamma^* \geq T$, we have already that $\mathbf{u} \in C^\infty(\mathbb{R}^3 \times (0, T))$, due to the arbitrariness of $\varepsilon \in (0, T)$. In the case when $\Gamma^* < T$, our strategy is to show that $u_3 \in L^3(\delta, \Gamma^*; L^9(\mathbb{R}^3))$. Then by the fact that

$$(1.15) \Rightarrow \partial_3 \mathbf{u} \in L^{5q/(2(2q-3))}(\delta, \Gamma^*; L^q(\mathbb{R}^3)) \subset L^{2q/(2q-3)}(\delta, \Gamma^*; L^q(\mathbb{R}^3)),$$

$$(1.16) \Rightarrow \partial_3 \mathbf{u} \in L^{11q/(7q-12)}(\delta, \Gamma^*; L^q(\mathbb{R}^3)) \subset L^{2q/(2q-3)}(\delta, \Gamma^*; L^q(\mathbb{R}^3)),$$

we may conclude the proof by invoking (1.17).

Case I. (1.15) holds. Multiplying the equation of u_3 in (1.1) by $|u_3|u_3$ and integrating over \mathbb{R}^3 , we obtain

$$(2.1) \quad \begin{aligned} \frac{1}{3} \frac{d}{dt} \| |u_3|^{3/2} \|_{L^2}^2 + \frac{4}{9} \| \nabla |u_3|^{3/2} \|_{L^2}^2 + \int_{\mathbb{R}^3} |u_3| \cdot |\nabla u_3|^2 dx \\ = - \int_{\mathbb{R}^3} \partial_3 \pi |u_3| u_3 dx \equiv I. \end{aligned}$$

We just need to estimate I :

$$\begin{aligned}
(2.2) \quad I &\leq \|\partial_3 \pi\|_{L^3} \|u_3\|_{L^3}^2 \quad (\text{Hölder inequality}) \\
&\leq C \|\mathbf{u}\|_{L^{3q/(q-3)}} \|\partial_3 \mathbf{u}\|_{L^q} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{4/3} \quad (\text{by (1.18)}) \\
&\leq C \|\mathbf{u}\|_{L^2}^{2(q-4)/(3q-2)} \|\mathbf{u}\|_{L^{3q}}^{(q+6)/(3q-2)} \|\partial_3 \mathbf{u}\|_{L^q} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{4/3} \\
&\quad (\text{interpolation inequality}) \\
&\leq C \|\partial_1 \mathbf{u}\|_{L^2}^{(q+6)/(3(3q-2))} \|\partial_2 \mathbf{u}\|_{L^2}^{(q+6)/(3(3q-2))} \|\partial_3 \mathbf{u}\|_{L^q}^{(q+6)/(3(3q-2))} \\
&\quad \times \|\partial_3 \mathbf{u}\|_{L^q} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{4/3} \\
&\quad (\text{by Definition 1.1 item 1, and (1.6)}) \\
&\leq C \|\nabla \mathbf{u}\|_{L^2}^{2(q+6)/(3(3q-2))} \|\partial_3 \mathbf{u}\|_{L^q}^{10q/(3(3q-2))} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{4/3} \\
&\leq C (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\partial_3 \mathbf{u}\|_{L^q}^{5q/(4q-6)}) \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{4/3} \quad (\text{Young inequality}).
\end{aligned}$$

Plugging (2.2) into (2.1), we find

$$\frac{d}{dt} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^2 \leq C (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\partial_3 \mathbf{u}\|_{L^q}^{5q/(2(2q-3))}) \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{4/3}.$$

Dividing both sides by $\|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{4/3}$ and integrating with respect to t , we deduce by assumption (1.15) that

$$\|u_3\|_{L^\infty(\delta, \Gamma^*; L^3(\mathbb{R}^3))} \leq C.$$

Integrating (2.1) then yields

$$\|u_3\|_{L^3(\delta, \Gamma^*; L^9(\mathbb{R}^3))} = \|u_3\|_{L^2(\delta, \Gamma^*; L^6(\mathbb{R}^3))}^{3/2} \leq C \|\nabla |u_3|^{3/2}\|_{L^2(\delta, \Gamma^*; L^2(\mathbb{R}^3))} \leq C,$$

as desired.

Case II. (1.16) holds. We should dominate I in another way:

$$\begin{aligned}
(2.3) \quad I &\leq C \int_{\mathbb{R}^3} |\pi| \cdot |\partial_3 u_3| \cdot |u_3| \, dx \\
&\leq C \|\pi\|_{L^{3q/(2q-3)}} \|\partial_3 u_3\|_{L^q} \|u_3\|_{L^3} \quad (\text{Hölder inequality}) \\
&\leq C \|\mathbf{u}\|_{L^{6q/(2q-3)}}^2 \|\partial_3 u_3\|_{L^q} \|u_3\|_{L^3} \quad (\text{by (1.18)}) \\
&\leq C \|\mathbf{u}\|_{L^2}^{2(2q-5)/(3q-2)} \|\mathbf{u}\|_{L^{3q}}^{2(q+3)/(3q-2)} \|\partial_3 u_3\|_{L^q} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{2/3} \\
&\quad (\text{interpolation inequality}) \\
&\leq C \|\partial_1 \mathbf{u}\|_{L^2}^{2(q+3)/(3(3q-2))} \|\partial_2 \mathbf{u}\|_{L^2}^{2(q+3)/(3(3q-2))} \|\partial_3 \mathbf{u}\|_{L^q}^{2(q+3)/(3(3q-2))} \\
&\quad \times \|\partial_3 u_3\|_{L^q} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{2/3} \\
&\quad (\text{by Definition 1.1 item 1, and (1.6)}) \\
&\leq C \|\nabla \mathbf{u}\|_{L^2}^{4(q+3)/(3(3q-2))} \|\partial_3 \mathbf{u}\|_{L^q}^{11q/(3(3q-2))} \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{2/3} \\
&\leq C (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\partial_3 \mathbf{u}\|_{L^q}^{11q/(7q-12)}) \|u_3\|_{L^2}^{3/2} \|u_3\|_{L^2}^{2/3} \quad (\text{Young inequality}).
\end{aligned}$$

Further, we may proceed as above to conclude the proof of Theorem 1.2 under condition (1.16).

References

- [1] *H. Beirão da Veiga*: A new regularity class for the Navier-Stokes equations in \mathbb{R}^n . *Chin. Ann. Math., Ser. B* *16* (1995), 407–412. [zbl](#) [MR](#)
- [2] *C. Cao*: Sufficient conditions for the regularity to the 3D Navier-Stokes equations. *Discrete Contin. Dyn. Syst.* *26* (2010), 1141–1151. [zbl](#) [MR](#) [doi](#)
- [3] *C. Cao, E. S. Titi*: Regularity criteria for the three-dimensional Navier-Stokes equations. *Indiana Univ. Math. J.* *57* (2008), 2643–2661. [zbl](#) [MR](#) [doi](#)
- [4] *J.-Y. Chemin, P. Zhang*: On the critical one component regularity for 3-D Navier-Stokes systems. *Ann. Sci. Éc. Norm. Supér. (4)* *49* (2016), 131–167. [zbl](#) [MR](#) [doi](#)
- [5] *J.-Y. Chemin, P. Zhang, Z. Zhang*: On the critical one component regularity for 3-D Navier-Stokes system: general case. *Arch. Ration. Mech. Anal.* *224* (2017), 871–905. [zbl](#) [MR](#) [doi](#)
- [6] *L. Escauriaza, G. A. Serëgin, V. Šverák*: $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness. *Russ. Math. Surv.* *58* (2003), 211–250; translation from *Usp. Mat. Nauk* *58* (2003), 3–44. [zbl](#) [MR](#) [doi](#)
- [7] *Z. Guo, M. CagGIO, Z. Skalák*: Regularity criteria for the Navier-Stokes equations based on one component of velocity. *Nonlinear Anal., Real World Appl.* *35* (2017), 379–396. [zbl](#) [MR](#) [doi](#)
- [8] *E. Hopf*: Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen. *Math. Nachr.* *4* (1951), 213–231. (In German.) [zbl](#) [MR](#) [doi](#)
- [9] *X. Jia, Y. Zhou*: Remarks on regularity criteria for the Navier-Stokes equations via one velocity component. *Nonlinear Anal., Real World Appl.* *15* (2014), 239–245. [zbl](#) [MR](#) [doi](#)
- [10] *I. Kukavica, M. Ziane*: One component regularity for the Navier-Stokes equations. *Nonlinearity* *19* (2006), 453–469. [zbl](#) [MR](#) [doi](#)
- [11] *I. Kukavica, M. Ziane*: Navier-Stokes equations with regularity in one direction. *J. Math. Phys.* *48* (2007), Article ID 065203, 10 pages. [zbl](#) [MR](#) [doi](#)
- [12] *J. Leray*: Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.* *63* (1934), 193–248. (In French.) [zbl](#) [MR](#) [doi](#)
- [13] *J. Neustupa, A. Novotný, P. Penel*: An interior regularity of a weak solution to the Navier-Stokes equations in dependence on one component of velocity. *Topics in Mathematical Fluid Mechanics. Quad. Mat.* *10*, Aracne, Rome, 2002, pp. 163–183. [zbl](#) [MR](#)
- [14] *P. Penel, M. Pokorný*: Some new regularity criteria for the Navier-Stokes equations containing gradient of the velocity. *Appl. Math., Praha* *49* (2004), 483–493. [zbl](#) [MR](#) [doi](#)
- [15] *P. Penel, M. Pokorný*: On anisotropic regularity criteria for the solutions to 3D Navier-Stokes equations. *J. Math. Fluid Mech.* *13* (2011), 341–353. [zbl](#) [MR](#) [doi](#)
- [16] *M. Pokorný*: On the result of He concerning the smoothness of solutions to the Navier-Stokes equations. *Electron. J. Differ. Equ.* *2003* (2003), Article No. 11, 8 pages. [zbl](#) [MR](#)
- [17] *G. Prodi*: Un teorema di unicITÀ per le equazioni di Navier-Stokes. *Ann. Mat. Pura Appl.* (4) *48* (1959), 173–182. (In Italian.) [zbl](#) [MR](#) [doi](#)
- [18] *J. C. Robinson, J. L. Rodrigo, W. Sadowski*: The Three-Dimensional Navier-Stokes Equations. *Classical Theory. Cambridge Studies in Advanced Mathematics* *157*, Cambridge University Press, Cambridge, 2016. [zbl](#) [MR](#) [doi](#)
- [19] *J. Serrin*: On the interior regularity of weak solutions of the Navier-Stokes equations. *Arch. Ration. Mech. Anal.* *9* (1962), 187–195. [zbl](#) [MR](#) [doi](#)
- [20] *Z. Skalák*: On the regularity of the solutions to the Navier-Stokes equations via the gradient of one velocity component. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* *104* (2014), 84–89. [zbl](#) [MR](#) [doi](#)

- [21] *R. Temam*: Navier-Stokes Equations. Theory and Numerical Analysis. AMS Chelsea Publishing, Providence, 2001. [zbl](#) [MR](#) [doi](#)
- [22] *Z. Zhang*: An almost Serrin-type regularity criterion for the Navier-Stokes equations involving the gradient of one velocity component. *Z. Angew. Math. Phys.* *66* (2015), 1707–1715. [zbl](#) [MR](#) [doi](#)
- [23] *Z. Zhang*: An improved regularity criterion for the Navier-Stokes equations in terms of one directional derivative of the velocity field. *Bull. Math. Sci.* *8* (2018), 33–47. [zbl](#) [MR](#) [doi](#)
- [24] *Y. Zhou*: A new regularity criterion for the Navier-Stokes equations in terms of the gradient of one velocity component. *Methods Appl. Anal.* *9* (2002), 563–578. [zbl](#) [MR](#) [doi](#)
- [25] *Y. Zhou*: A new regularity criterion for weak solutions to the Navier-Stokes equations. *J. Math. Pures Appl.* (9) *84* (2005), 1496–1514. [zbl](#) [MR](#) [doi](#)
- [26] *Y. Zhou, M. Pokorný*: On a regularity criterion for the Navier-Stokes equations involving gradient of one velocity component. *J. Math. Phys.* *50* (2009), Article ID 123514, 11 pages. [zbl](#) [MR](#) [doi](#)
- [27] *Y. Zhou, M. Pokorný*: On the regularity of the solutions of the Navier-Stokes equations via one velocity component. *Nonlinearity* *23* (2010), 1097–1107. [zbl](#) [MR](#) [doi](#)

Authors' addresses: *Zujin Zhang, Weijun Yuan*, School of Mathematics and Computer Sciences, Gannan Normal University, Ganzhou 341000, Jiangxi, P. R. China, e-mail: zhangzujin361@163.com, 1453540745@qq.com; *Yong Zhou*, School of Mathematics (Zhuhai), Sun Yat-sen University, Zhuhai, Guangdong 519082, P. R. China; Department of Mathematics, Zhejiang Normal University, Jinhua 321004, Zhejiang, P. R. China, e-mail: zhouyong3@mail.sysu.edu.cn.