

ASYMPTOTIC LOWER BOUNDS FOR EIGENVALUES OF THE  
STEKLOV EIGENVALUE PROBLEM WITH VARIABLE  
COEFFICIENTS

YU ZHANG, HAI BI, YIDU YANG, Guiyang

Received April 26, 2019. Published online September 9, 2020.

*Abstract.* In this paper, using a new correction to the Crouzeix-Raviart finite element eigenvalue approximations, we obtain asymptotic lower bounds of eigenvalues for the Steklov eigenvalue problem with variable coefficients on  $d$ -dimensional domains ( $d = 2, 3$ ). In addition, we prove that the corrected eigenvalues converge to the exact ones from below. The new result removes the conditions of eigenfunction being singular and eigenvalue being large enough, which are usually required in the existing arguments about asymptotic lower bounds. Further, we prove that the corrected eigenvalues still maintain the same convergence order as uncorrected eigenvalues. Finally, numerical experiments validate our theoretical results.

*Keywords:* correction; Steklov eigenvalue problem; Crouzeix-Raviart finite element; asymptotic lower bounds; convergence order

*MSC 2020:* 65N25, 65N30

## 1. INTRODUCTION

It is an important topic to obtain upper and lower bounds for eigenvalues. As we all know, thanks to the minimum-maximum principle, it is easy to obtain guaranteed upper bounds of eigenvalues by conforming finite element methods. Naturally, attention has been paid to finding lower bounds of eigenvalues by nonconforming finite elements, such as the rotated bilinear ( $Q_1^{\text{rot}}$ ) finite element [21], [23], [15], [19], the extension of  $Q_1^{\text{rot}}$  finite element [21], [18], [19], [16], the enriched Crouzeix-Raviart (ECR) finite element [15], [16], [19], [22], [25], the Wilson finite element [21], [36],

---

This work is supported by the Young Scientific and Technical Talents Development of Education Department of Guizhou Province (KY [2018]153), the National Natural Science Foundation of China (Grant No. 11561014 and No. 11761022).

the Morley element [7], [16], [32], etc. Especially, a lot of work has been done on asymptotic lower bounds for eigenvalues based on the Crouzeix-Raviart (CR) finite element approximations (see, e.g., [2], [23], [34], [16], [30], [19], [31] and the citations therein). Asymptotic lower bounds require that the mesh size is small enough. Recently, finding guaranteed lower bounds has become an attractive topic, which has no requirement for the mesh size. There are also some researches on finding guaranteed lower bounds for eigenvalues based on the CR finite element (see, e.g., [8], [24], [17], [29], [35]).

In this paper, we discuss asymptotic lower bounds for eigenvalues of the Steklov eigenvalue problem with variable coefficients

$$(1.1) \quad \begin{cases} -\operatorname{div}(\alpha \nabla u) + \beta u = 0 & \text{in } \Omega, \\ \alpha \frac{\partial u}{\partial \nu} = \lambda u & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega \subset \mathbb{R}^d$  ( $d = 2, 3$ ) is a bounded polygonal or polyhedral domain and  $\partial u / \partial \nu$  is the outward normal derivative on  $\partial\Omega$ . Symbols  $\nabla$  and  $\operatorname{div}$  denote the gradient and the divergence operators, respectively,  $\beta = \beta(x) \in L^\infty(\Omega)$  has a positive lower bound,  $\alpha = \alpha(x) \in W^{1,\infty}(\Omega)$ , and  $\alpha_0 \leq \alpha(x)$  for a constant  $\alpha_0 > 0$ .

Among the above references for the Steklov eigenvalue problem with constant coefficients, [19], [31], [22] and [35] discuss asymptotic and guaranteed lower eigenvalue bounds, respectively. The paper [19] states that the CR finite element produces asymptotic lower bounds for eigenvalues in the case of a singular eigenfunction. And [19], [31] also prove that property of asymptotic lower bounds in the case of a non-singular eigenfunction but under an additional condition that the eigenvalue is large enough. The paper [35] obtains guaranteed lower bounds for eigenvalue by correcting the CR finite element eigenvalue approximations, but convergence order of the corrected eigenvalues cannot achieve that of the uncorrected eigenvalues.

Based on the above work, we further discuss asymptotic lower bounds of eigenvalues for the Steklov eigenvalue problem with variable coefficients. We introduce a new correction formula (3.5) to the CR finite element eigenvalue approximations  $\lambda_h$  and obtain the corrected eigenvalues  $\lambda_h^c$ . Our work has the following features:

(1) For shape-regular meshes including quasi-uniform meshes and adaptive meshes with local refinement, when mesh diameter  $h$  is sufficiently small and  $\|u - u_h\| \geq Ch^{1+\varepsilon_0}$  (for some  $\varepsilon_0 > 0$ ), we prove the conclusion

$$\lambda \geq \lambda_h^c$$

in Theorem 3.1, which shows that the corrected eigenvalues are asymptotic lower bounds of the exact ones. The new result removes the conditions of eigenfunction

being singular and eigenvalue being large enough (see Section 3 for details), which are usually required in the existing arguments about asymptotic lower bounds.

(2) The result in Theorem 3.2 implies that the corrected eigenvalues converge to the exact ones without the loss of convergence order, i.e., convergence order of the corrected eigenvalues is still the same as that of the uncorrected eigenvalues.

(3) For  $d$ -dimensional domains ( $d = 2, 3$ ), we implement numerical experiments in Section 4. Numerical results coincide in the theoretical analysis. We are particularly pleased that the correction takes very little time.

In [19], [22], it has been obtained that the ECR finite element can produce asymptotic lower bounds of eigenvalues for the Steklov eigenvalue problem with a constant coefficient whether the eigenfunctions are singular or not. However, up to now, it has not been proved that the ECR element produces asymptotic lower eigenvalue bounds for the Steklov eigenvalue problem with variable coefficients. It should be pointed out that the correction method and theoretical analysis in this paper are also valid for the ECR finite element (see Remark 3.3 in Section 3).

As for the basic theory of the finite element and spectral approximation, we refer to [3], [4], [26], [6], [1], and [5]. Throughout this paper,  $C$  denotes a generic positive constant independent of the mesh size, which may not be the same at each occurrence.

## 2. PRELIMINARY

Let  $H^m(\Omega)$  denote the Sobolev space with the real order  $m$  on  $\Omega$ . Let  $\|\cdot\|_{m,\Omega}$  and  $|\cdot|_{m,\Omega}$  be the norm and seminorm on  $H^m(\Omega)$ , respectively, and  $H^0(\Omega) = L^2(\Omega)$ . Furthermore,  $H^m(\partial\Omega)$  denotes the Sobolev space with the real order  $m$  on  $\partial\Omega$ ,  $\|\cdot\|_{m,\partial\Omega}$  is the norm on  $H^m(\partial\Omega)$  and  $H^0(\partial\Omega) = L^2(\partial\Omega)$ .

The weak form of (1.1) can be written as: find  $(\lambda, u) \in \mathbb{R} \times H^1(\Omega)$  with  $\|u\|_{0,\partial\Omega} = 1$  such that

$$(2.1) \quad a(u, v) = \lambda b(u, v) \quad \forall v \in H^1(\Omega),$$

where

$$(2.2) \quad a(u, v) = \int_{\Omega} (\alpha \nabla u \cdot \nabla v + \beta uv) \, dx,$$

$$(2.3) \quad b(u, v) = \int_{\partial\Omega} uv \, ds.$$

Let  $\pi_h = \{\kappa\}$  be a regular partition of  $\Omega$  with the mesh diameter  $h = \max\{h_{\kappa}\}$ , where  $h_{\kappa}$  is the diameter of element  $\kappa$ . Let  $\varepsilon_h$  be the set of  $(d-1)$ -dimensional faces of  $\pi_h$ . We denote by  $|\kappa|$  the measure of the element  $\kappa$ .

We consider the CR finite element space  $V_h$ , proposed by Crouzeix and Raviart [12], as

$$V_h = \{v \in L^2(\Omega) : v|_\kappa \in P_1(\kappa), v \text{ is continuous at the barycenters of the } (d-1)\text{-dimensional faces of } \kappa \text{ for all } \kappa \in \pi_h\}.$$

Put  $\|v\|_h = \left(\sum_{\kappa \in \pi_h} \|v\|_{1,\kappa}^2\right)^{1/2}$ , then  $\|v\|_h$  means the norm on  $V_h$ . To construct the CR finite element approximation of (2.1) means to find  $(\lambda_h, u_h) \in \mathbb{R} \times V_h$  with  $\|u_h\|_{0,\partial\Omega} = 1$  such that

$$(2.4) \quad a_h(u_h, v) = \lambda_h b(u_h, v) \quad \forall v \in V_h,$$

where

$$(2.5) \quad a_h(u_h, v) = \sum_{\kappa \in \pi_h} \int_{\kappa} (\alpha \nabla u_h \cdot \nabla v + \beta u_h v) dx.$$

From Theorem 4 in [27] and Remark 2.1 in [14], we have the following regularity result.

**Regularity:** Assume that  $\varphi$  is the solution of the source problem associated with (2.1) with the right-hand side  $f$ .

▷ In the case of  $\Omega \subset \mathbb{R}^2$ , if  $f \in L^2(\partial\Omega)$ , there exist constants  $r$  and  $C_r > 0$  such that  $\varphi \in H^{1+r}(\Omega)$  and

$$\|\varphi\|_{1+r/2} \leq C_r \|f\|_{0,\partial\Omega};$$

if  $f \in H^{1/2}(\partial\Omega)$ , then  $\varphi \in H^{1+r}(\Omega)$  and

$$\|\varphi\|_{1+r} \leq C_r \|f\|_{1/2,\partial\Omega},$$

here  $r = 1$  when the largest inner angle  $\theta$  of  $\Omega$  satisfies  $\theta < \pi$ , and  $r < \pi/\theta$  which can be arbitrarily close to  $\pi/\theta$  when  $\theta > \pi$ .

▷ In the case of  $\Omega \subset \mathbb{R}^3$ , if  $f \in L^2(\partial\Omega)$ , there exist constants  $r \in (0, \frac{1}{2})$  and  $C_r > 0$  such that  $\varphi \in H^{1+r}(\Omega)$  and

$$\|\varphi\|_{1+r} \leq C_r \|f\|_{0,\partial\Omega},$$

where  $C_r$  is the regularity constant independent of  $f$ .

**Lemma 2.1.** *Let  $(\lambda_h, u_h)$  be the  $j$ th eigenpair of (2.4) and  $\lambda$  be the  $j$ th eigenvalue of (2.1). Then there exists an eigenfunction  $u$  corresponding to  $\lambda$  and when  $u \in$*

$H^{1+t}(\Omega)$  and  $h$  is sufficiently small, it holds

$$(2.6) \quad \|u_h - u\|_h \leq Ch^t,$$

$$(2.7) \quad |\lambda_h - \lambda| \leq Ch^{2t},$$

$$(2.8) \quad \|u - u_h\|_{0,\partial\Omega} \leq Ch^s \|u - u_h\|_h,$$

where  $r \leq t \leq 1$ ,  $s = r/2$  if  $\Omega \subset \mathbb{R}^2$  and  $s = r$  if  $\Omega \subset \mathbb{R}^3$ . The eigenfunction  $u$  is called singular when  $t < 1$ .

*Proof.* By using standard arguments in nonconforming finite element error estimates, (2.6) and (2.7) can be proved directly. The two estimates have also been given by Theorem 2.2 in [13] and (2.6) in [14]. Now we prove (2.8).

In order to prove the error estimate, we need to define the solution operator  $A: L^2(\partial\Omega) \rightarrow H^1(\Omega)$  associated with the source problem of (2.1) by

$$a(Af, v) = b(f, v) \quad \forall v \in H^1(\Omega)$$

and the operator  $T: L^2(\partial\Omega) \rightarrow L^2(\partial\Omega)$  by

$$Tf = (Af)',$$

where the prime denotes the restriction to  $\partial\Omega$ , namely  $Tf = Af|_{\partial\Omega}$ .

Analogously, we can define the discrete versions  $A_h$  and  $T_h$  corresponding to  $A$  and  $T$ , respectively. Define  $A_h: L^2(\partial\Omega) \rightarrow V_h$  by

$$a_h(A_h f, v) = b(f, v) \quad \forall v \in V_h$$

and the operator  $T_h: L^2(\partial\Omega) \rightarrow L^2(\partial\Omega)$  by

$$T_h f = (A_h f)'.$$

By the Nitsche technique (see also (2.13) in [30]), we derive

$$\|Tu - T_h u\|_{0,\partial\Omega} = \|Au - A_h u\|_{0,\partial\Omega} \leq Ch^s \|Au - A_h u\|_h.$$

From (2.7) and (2.8) in [34] (see also Lemma 3.1 in [19]), we have

$$\begin{aligned} \|u - u_h\|_h &= \lambda \|Au - A_h u\|_h + R, \\ \|u - u_h\|_{0,\partial\Omega} &\leq C \|Tu - T_h u\|_{0,\partial\Omega}, \end{aligned}$$

where  $|R| \leq C \|Tu - T_h u\|_{0,\partial\Omega}$ .

The estimate (2.8) is a direct consequence of the above three relations.  $\square$

Define the Crouzeix-Raviart interpolation operator  $I_h : H^1(\Omega) \rightarrow V_h$  by

$$(2.9) \quad \int_e I_h u \, ds = \int_e u \, ds \quad \forall e \in \varepsilon_h, \quad u \in H^1(\Omega).$$

Note that the interpolation operator  $I_h$  has an important orthogonality property (see the equality (2.9) in [2]): for each element  $\kappa \in \pi_h$ , it is

$$(2.10) \quad \int_{\kappa} \nabla(u - I_h u) \cdot \nabla v_h \, dx = \int_{\partial\kappa} (u - I_h u) \nabla v_h \cdot \nu \, ds = 0 \quad \forall v_h \in V_h.$$

The estimation of constants in the Poincaré and trace inequalities is a concern of academe (see, e.g., [28], [9], [7], [10], [20], [35], [24] and therein). From Theorem 4.2 in [24], we have following Lemma 2.2.

**Lemma 2.2.** *For any element  $\kappa$ , the conclusion*

$$(2.11) \quad \|u - I_h u\|_{0,\kappa} \leq C_{h_\kappa} |u - I_h u|_{1,\kappa} \quad \forall u \in H^1(\kappa),$$

is valid where

- ▷  $C_{h_\kappa} = 0.1893h_\kappa$  for a triangle element  $\kappa$  in  $\mathbb{R}^2$ ,
- ▷  $C_{h_\kappa} = 0.3804h_\kappa$  for a tetrahedron element  $\kappa$  in  $\mathbb{R}^3$ .

Consider any element  $\kappa$  with the vertices  $P_1, P_2, \dots, P_{d+1}$ . The edge/face opposite to the vertex  $P_{d+1}$  is denoted by  $e$ . The measure of  $e$  is  $|e|$ . Let  $H_\kappa$  be the height of element  $\kappa$  with respect to  $e$ . It is easy to see that

$$H_\kappa = \frac{d|\kappa|}{|e|}.$$

Thanks to Lemma 2 of [7] and Theorem 3.3 of [35], we have following Lemma 2.3.

**Lemma 2.3.** *For a given element  $\kappa$ , it is*

$$(2.12) \quad \|u - I_h u\|_{0,e} \leq C_{h_e} |u - I_h u|_{1,\kappa} \quad \forall u \in H^1(\kappa),$$

where

- ▷  $C_{h_e} = 0.6711 \frac{h_\kappa}{\sqrt{H_\kappa}}$  for a triangle element  $\kappa$  in  $\mathbb{R}^2$ ,
- ▷  $C_{h_e} = 1.0932 \frac{h_\kappa}{\sqrt{H_\kappa}}$  for a tetrahedron element  $\kappa$  in  $\mathbb{R}^3$ .

Proof. The proof can be found in Theorem 3.3 of [35]. For convenience of reading in case of  $d = 3$ , we present the proof here.

For any  $v \in H^1(\kappa)$  and any point  $x = (x_1, x_2, x_3)$  in  $\kappa$ , we have

$$(2.13) \quad \int_{\kappa} ((x_1, x_2, x_3) - P_4) \cdot \nabla(v^2) \, dx = \int_{\partial\kappa} ((x_1, x_2, x_3) - P_4) \cdot \mathbf{n} v^2 \, ds - \int_{\kappa} 3v^2 \, dx$$

from the Green formula. We deduce

$$(2.14) \quad \begin{aligned} & ((x_1, x_2, x_3) - P_4) \cdot \mathbf{n} \\ &= \begin{cases} 0 & \text{for any } x \text{ on the faces } P_1P_2P_4, P_1P_3P_4, \text{ and } P_2P_3P_4, \\ \frac{3|\kappa|}{|e|} & \text{for any } x \text{ on the face } P_1P_2P_3. \end{cases} \end{aligned}$$

Substituting (2.14) into (2.13), we obtain

$$(2.15) \quad \begin{aligned} \frac{3|\kappa|}{|e|} \int_e v^2 \, ds &= \int_{\kappa} 3v^2 \, dx + \int_{\kappa} ((x_1, x_2, x_3) - P_4) \cdot \nabla(v^2) \, dx \\ &\leq 3 \int_{\kappa} v^2 \, dx + \int_{\kappa} |(x_1, x_2, x_3) - P_4| |\nabla(v^2)| \, dx \\ &\leq 3 \int_{\kappa} v^2 \, dx + 2h_{\kappa} \int_{\kappa} |v| |\nabla v| \, dx \\ &\leq 3 \|v\|_{0,\kappa}^2 + 2h_{\kappa} \|v\|_{0,\kappa} \|\nabla v\|_{0,\kappa}. \end{aligned}$$

Taking  $v = u - I_h u$  and applying the estimate (2.11), we deduce

$$\|u - I_h u\|_{0,e}^2 \leq \frac{|e|}{3|\kappa|} (3C_{h_{\kappa}}^2 + 2h_{\kappa} C_{h_{\kappa}}) \|u - I_h u\|_{1,\kappa}^2,$$

which implies that (2.12) is valid when  $\Omega \subset \mathbb{R}^3$ . □

### 3. THE ASYMPTOTIC LOWER BOUNDS PROPERTY OF CORRECTED EIGENVALUES

For the problem (1.1), thanks to the minimum-maximum principle, it is easy to obtain guaranteed upper bounds for eigenvalues by conforming finite element methods. From [19], we know that the CR finite element method gives asymptotic lower bounds for eigenvalues when the corresponding eigenfunctions are singular or the eigenvalues are large enough. In this section, we introduce a correction for eigenvalues of the problem (1.1) and we prove that the corrected eigenvalues converge to the exact ones from below. The conclusion holds without the conditions that

eigenfunction is singular and eigenvalue is large enough. First we prove the following inequality (3.1) and Lemma 3.1.

Using (2.10), we have

$$\int_{\kappa} \nabla(u - I_h u) \cdot \nabla(u - I_h u) \, dx = \int_{\kappa} \nabla(u - I_h u) \cdot \nabla(u - u_h) \, dx \leq |u - I_h u|_{1,\kappa} |u - u_h|_{1,\kappa},$$

so then

$$(3.1) \quad |u - I_h u|_{1,\kappa} \leq |u - u_h|_{1,\kappa}.$$

The identity in following Lemma 3.1 is an equivalent form of the identity (4.1) in [19], which is a generalization of the identities (2.12) in [2] and (2.3) in [36].

**Lemma 3.1.** *Let  $(\lambda, u)$  and  $(\lambda_h, u_h)$  be eigenpairs of (2.1) and (2.4), respectively. Then the following identity is valid:*

$$(3.2) \quad \begin{aligned} \lambda - \lambda_h &= a_h(u - u_h, u - u_h) - \lambda_h b(u - u_h, u - u_h) \\ &\quad - 2a_h(I_h u - u, u_h) - 2\lambda_h b(u - I_h u, u_h). \end{aligned}$$

*Proof.* From  $\|u\|_{0,\partial\Omega} = 1 = \|u_h\|_{0,\partial\Omega}$ , we get

$$a_h(u, u) = \lambda, \quad a_h(u_h, u_h) = \lambda_h.$$

Therefore,

$$(3.3) \quad \begin{aligned} \lambda - \lambda_h &= a_h(u, u) + a_h(u_h, u_h) - 2a_h(u_h, u_h) \\ &= a_h(u, u) + a_h(u_h, u_h) - 2a_h(u, u_h) + 2a_h(u - u_h, u_h) \\ &= a_h(u - u_h, u - u_h) + 2a_h(u - u_h, u_h). \end{aligned}$$

From  $b(I_h u - u_h, u_h) = b(I_h u - u, u_h) + b(u - u_h, u_h - \frac{1}{2}u + \frac{1}{2}u)$ , we obtain

$$\lambda_h b(I_h u - u_h, u_h) = \lambda_h b(I_h u - u, u_h) - \frac{1}{2}\lambda_h b(u - u_h, u - u_h),$$

which together with (2.4) yields

$$(3.4) \quad \begin{aligned} a_h(u - u_h, u_h) &= a_h(u - I_h u, u_h) + a_h(I_h u - u_h, u_h) \\ &= a_h(u - I_h u, u_h) + \lambda_h b(I_h u - u_h, u_h) \\ &= a_h(u - I_h u, u_h) + \lambda_h b(I_h u - u, u_h) - \frac{1}{2}\lambda_h b(u - u_h, u - u_h). \end{aligned}$$

Substituting (3.4) into (3.3), we get (3.2). □



Now we give the correction formula (3.5). In addition, we prove that the correction provides asymptotic lower bounds for eigenvalues of the problem (2.1).

Denote by  $I_0$  the piecewise constant interpolation operator on  $\Omega$ . Let  $(\lambda, u)$  be an eigenpair of (2.1) and  $(\lambda_h, u_h)$  be the corresponding CR finite element approximations. We introduce the following formula to correct the CR finite element approximations  $\lambda_h$ :

$$(3.5) \quad \lambda_h^c = \frac{\lambda_h}{1 + M/\lambda_h},$$

where

$$(3.6) \quad M = \frac{\delta}{\alpha_0} \sum_{\kappa \in \pi_h} (\|(\alpha - I_0\alpha)\nabla u_h\|_{0,\kappa} + Ch_\kappa \|\beta u_h\|_{0,\kappa})^2$$

and  $\delta > 1$  is any given constant.

By the interpolation error estimate, we know that

$$(3.7) \quad \|\alpha - I_0\alpha\|_{0,\infty,\kappa} \leq Ch_\kappa \|\alpha\|_{1,\infty,\kappa}.$$

Noting that  $Ch_\kappa = 0.1893h_\kappa$ , we derive

$$(3.8) \quad 0 \leq M \leq Ch^2.$$

In practical computation, we cannot guarantee that  $\lambda_h$  are lower bounds of  $\lambda$  if we are not sure that the eigenfunctions are singular or the eigenvalues are large enough. Now we prove that the corrected eigenvalues  $\lambda_h^c$  are asymptotic lower bounds of the exact ones, which holds without the conditions of singularity and large eigenvalues.

**Theorem 3.1.** *Let  $\lambda_h^c$  be a corrected eigenvalue obtained by (3.5). Assume that the conditions of Lemma 2.1 and  $\|u - u_h\|_h \geq Ch^{1+\varepsilon_0}$  ( $\varepsilon_0 = \min\{\frac{1}{4}, \frac{1}{2}r\}$ ) hold. Then, if  $h$  is sufficiently small, we have*

$$(3.9) \quad \lambda \geq \lambda_h^c.$$

**Proof.** We now estimate each of the four terms on the right-hand side of (3.2). Since  $\alpha \geq \alpha_0$ , the first term

$$(3.10) \quad a_h(u - u_h, u - u_h) \geq \sum_{\kappa \in \pi_h} \left( \alpha_0 |u - u_h|_{1,\kappa}^2 + \int_\kappa \beta (u - u_h)^2 dx \right).$$

From (2.8), the second term

$$(3.11) \quad \lambda_h b(u - u_h, u - u_h) = \lambda_h \|u - u_h\|_{0,\partial\Omega}^2 \leq Ch^{2s} \|u - u_h\|_h^2.$$

Now we estimate the third term. From (2.10), we have

$$(3.12) \quad \begin{aligned} a_h(I_h u - u, u_h) &= \sum_{\kappa \in \pi_h} \int_{\kappa} ((\alpha - I_0 \alpha) \nabla(I_h u - u) \cdot \nabla u_h \\ &\quad + I_0 \alpha \nabla(I_h u - u) \cdot \nabla u_h + \beta(I_h u - u) u_h) \, dx \\ &= \sum_{\kappa \in \pi_h} \int_{\kappa} ((\alpha - I_0 \alpha) \nabla(I_h u - u) \cdot \nabla u_h + \beta(I_h u - u) u_h) \, dx. \end{aligned}$$

Applying Schwarz's inequality and (2.11) to the above equality and combining it with (3.1), we deduce that

$$\begin{aligned} a_h(I_h u - u, u_h) &\leq \sum_{\kappa \in \pi_h} (|u - I_h u|_{1,\kappa} \|(\alpha - I_0 \alpha) \nabla u_h\|_{0,\kappa} + \|u - I_h u\|_{0,\kappa} \|\beta u_h\|_{0,\kappa}) \\ &\leq \sum_{\kappa \in \pi_h} |u - I_h u|_{1,\kappa} (\|(\alpha - I_0 \alpha) \nabla u_h\|_{0,\kappa} + C_{h_\kappa} \|\beta u_h\|_{0,\kappa}) \\ &\leq \sum_{\kappa \in \pi_h} |u - u_h|_{1,\kappa} (\|(\alpha - I_0 \alpha) \nabla u_h\|_{0,\kappa} + C_{h_\kappa} \|\beta u_h\|_{0,\kappa}), \end{aligned}$$

which together with Young's inequality yields

$$(3.13) \quad \begin{aligned} 2a_h(I_h u - u, u_h) &\leq \frac{\alpha_0}{\delta} \sum_{\kappa \in \pi_h} |u - u_h|_{1,\kappa}^2 \\ &\quad + \frac{\delta}{\alpha_0} \sum_{\kappa \in \pi_h} (\|(\alpha - I_0 \alpha) \nabla u_h\|_{0,\kappa} + C_{h_\kappa} \|\beta u_h\|_{0,\kappa})^2. \end{aligned}$$

It remains to estimate the last term. For the later proof, we introduce the piecewise constant interpolation operator  $I_0^b$  on  $\partial\Omega$ . Using (2.9), Schwarz's inequality, (2.12),  $\|u - I_0^b u\|_{0,e} \leq Ch^{\min\{1, 1/2+r\}} \|u\|_{H^{1/2+r}(e)}$ , the trace inequality and (3.1), we get

$$\begin{aligned} b(u - I_h u, u) &= \sum_{e \in \varepsilon_h \cap \partial\Omega} \int_e ((u - I_h u)(u - I_0^b u) + (u - I_h u) I_0^b u) \, ds \\ &\leq \sum_{e \in \varepsilon_h \cap \partial\Omega} \|u - I_h u\|_{0,e} \|u - I_0^b u\|_{0,e} \\ &\leq Ch^{\min\{1, 1/2+r\}} \left( \sum_{\substack{\kappa \in \pi_h, \\ e \in \partial\kappa \cap \partial\Omega}} C_{h_e}^2 |u - I_h u|_{1,\kappa}^2 \right)^{1/2} \\ &\leq Ch^{1+2\varepsilon_0} \left( \sum_{\kappa \in \pi_h} |u - u_h|_{1,\kappa}^2 \right)^{1/2}, \end{aligned}$$

where  $\varepsilon_0 = \min\{\frac{1}{4}, \frac{1}{2}r\}$ . Combining it with  $\|u - u_h\|_h \geq Ch^{1+\varepsilon_0}$ , we derive that

$$(3.14) \quad b(u - I_h u, u) \leq Ch^{\varepsilon_0} \|u - u_h\|_h^2.$$

From Schwarz's inequality, (2.12), (2.8) and (3.1), we conclude that

$$(3.15) \quad \begin{aligned} b(u - I_h u, u_h - u) &\leq \sum_{e \in \varepsilon_h \cap \partial\Omega} \|u - I_h u\|_{0,e} \|u_h - u\|_{0,e} \\ &\leq Ch^{1/2} \left( \sum_{\kappa \in \pi_h} |u - I_h u|_{1,\kappa}^2 \right)^{1/2} h^s \|u_h - u\|_h \\ &\leq Ch^{1/2+s} \|u_h - u\|_h^2. \end{aligned}$$

Combining (3.14) and (3.15), we deduce

$$(3.16) \quad 2\lambda_h b(u - I_h u, u_h) \leq Ch^{\varepsilon_0} \|u_h - u\|_h^2.$$

Substituting (3.10), (3.11), (3.13), (3.16) into (3.2), we obtain

$$(3.17) \quad \begin{aligned} \lambda - \lambda_h &\geq \left(1 - \frac{1}{\delta}\right) \alpha_0 \sum_{\kappa \in \pi_h} |u - u_h|_{1,\kappa}^2 \\ &\quad + \sum_{\kappa \in \pi_h} \int_{\kappa} \beta (u - u_h)^2 dx - Ch^{2s} \|u - u_h\|_h^2 \\ &\quad - \frac{\delta}{\alpha_0} \sum_{\kappa \in \pi_h} (\|(\alpha - I_0 \alpha) \nabla u_h\|_{0,\kappa} + C_{h,\kappa} \|\beta u_h\|_{0,\kappa})^2 - Ch^{\varepsilon_0} \|u_h - u\|_h^2. \end{aligned}$$

From the definition of  $M$ , we have

$$(3.18) \quad \begin{aligned} \left(1 + \frac{1}{\lambda_h} M\right) \lambda - \lambda_h &\geq \left(1 - \frac{1}{\delta}\right) \alpha_0 \sum_{\kappa \in \pi_h} |u - u_h|_{1,\kappa}^2 + \sum_{\kappa \in \pi_h} \int_{\kappa} \beta (u - u_h)^2 dx \\ &\quad - Ch^{2s} \|u - u_h\|_h^2 - Ch^{\varepsilon_0} \|u_h - u\|_h^2 - \frac{\lambda_h - \lambda}{\lambda_h} M. \end{aligned}$$

It is obvious that, when  $h$  is sufficiently small, the third and the fourth terms on the right-hand side of (3.18) are infinitesimals of higher order compared with the sum of the first two terms. From (3.8), (2.7) and  $\|u - u_h\|_h \geq Ch^{1+\varepsilon_0}$ , we get

$$\left| \frac{\lambda_h - \lambda}{\lambda_h} M \right| \leq Ch^{2+2t} \leq Ch^t h^{2+t} \leq Ch^t \|u - u_h\|_h^2,$$

which is a quantity of high order. Hence, the sign of the right-hand side of (3.18) is determined by the sum of the first two terms, i.e.

$$\left(1 + \frac{1}{\lambda_h} M\right) \lambda - \lambda_h \geq 0.$$

From (3.5), we know that (3.9) is valid. The proof is completed.  $\square$

**Remark 3.1.** A condition on the lower bound for  $\|u - u_h\|_h$  is necessary in Theorem 3.1, otherwise the proof does not work. The condition  $\|u - u_h\|_h \geq Ch^r$  has been used in Theorem 2.3 of [2]. It is valid on quasi-uniform meshes but not on adaptive meshes with local refinement. Therefore, in order to make the conclusion of Theorem 3.1 hold on shape-regular meshes including quasi-uniform meshes and adaptive meshes with local refinement, we use the condition  $\|u - u_h\|_h \geq Ch^{1+\varepsilon_0}$  ( $\varepsilon_0 = \min\{\frac{1}{4}, \frac{1}{2}r\}$ ) rather than  $\|u - u_h\|_h \geq Ch^r$ . There are some papers on the adaptive algorithm that have discussed the rationality of this type of hypothesis (see, e.g., (2.33) and Remark 2.1 in [33]).

The following theorem shows that  $\lambda_h^c$  converge to  $\lambda$  and maintain the same convergence order as  $\lambda_h$ .

**Theorem 3.2.** *Let  $(\lambda, u)$  and  $(\lambda_h, u_h)$  be eigenpairs of (2.1) and (2.4), respectively. If  $\lambda_h^c$  is a corrected eigenvalue obtained by (3.5), then we have*

$$(3.19) \quad \lambda - \lambda_h^c = \lambda - \lambda_h + \frac{\lambda_h M}{\lambda_h + M},$$

where  $|M| \leq Ch^2$ .

**Proof.** From (3.6), we have

$$\lambda - \lambda_h^c = \lambda - \lambda_h + \lambda_h - \frac{\lambda_h}{1 + M/\lambda_h} = \lambda - \lambda_h + \frac{\lambda_h M}{\lambda_h + M}.$$

The proof is completed. □

**Remark 3.2.** From (3.17) and the definition of  $M$ , we can also get another correction formula

$$\lambda_h^N = \lambda_h - M.$$

Here  $\lambda_h^N$  is still an asymptotic lower bound for the eigenvalue. In addition, it can be obtained directly that

$$\lambda - \lambda_h^N = \lambda - \lambda_h + M,$$

which indicates that  $\lambda_h^N$  converge to  $\lambda$  and maintain the same convergence order as  $\lambda_h$ . However, from the inequality

$$(\lambda - \lambda_h^c) - (\lambda - \lambda_h^N) = \frac{\lambda_h M}{\lambda_h + M} - M = -\frac{M^2}{\lambda_h + M} \leq 0,$$

we know that the error of  $\lambda_h^N$  is larger than  $\lambda_h^c$  when the mesh size is small enough.

**Remark 3.3** (The correction to the ECR finite element eigenvalue approximations). Let  $(\lambda_h, u_h)$  be an approximate eigenpair of (2.1) obtained by the ECR element and  $\beta \in W^{1,\infty}(\Omega)$ . Assume that the condition  $\|u - u_h\|_h \geq Ch^{1+\varepsilon_0}$  holds. Following closely the arguments used to prove Theorem 3.1, if  $h$  is sufficiently small, we can also obtain a similar correction

$$\lambda_h^c = \frac{\lambda_h}{1 + \delta \lambda_h^{-1} \alpha_0^{-1} \sum_{\kappa \in \pi_h} \|(\alpha - I_o \alpha) \nabla u_h\|_{0,\kappa}^2}$$

such that

$$\lambda \geq \lambda_h^c$$

and  $\lambda_h^c$  maintain the same convergence order as  $\lambda_h$ .

Actually, throughout the proof, we just need to replace the second term in (3.12) with the term

$$\begin{aligned} \sum_{\kappa \in \pi_h} \int_{\kappa} \beta(u - I_h u) u_h \, dx &= \sum_{\kappa \in \pi_h} \int_{\kappa} (u - I_h u) (\beta u_h - I_0(\beta u_h)) \, dx \\ &\leq C \sum_{\kappa \in \pi_h} h_{\kappa}^2 |u - I_h u|_{1,\kappa} \|\beta u_h\|_{1,\kappa}, \end{aligned}$$

which is a quantity of high order. Then we can obtain the desired.

#### 4. NUMERICAL EXPERIMENTS

In this section, to validate the theoretical results of this paper, we apply the correction (3.5) to (1.1) on the domain  $\Omega$ . The discrete eigenvalue problems are solved in MATLAB 2018b on an Lenovo IdeaPad PC with 1.8 GHz CPU and 8 GB RAM. Our code was compiled under the iFEM package [11]. In order to investigate the error, we use the approximate eigenvalues given by extrapolation method as the reference value. The following notations are adopted in tables and figures.

$h_0$ : The diameter of  $\Omega$ .

$h$ : The diameter of meshes.

$\lambda_j$ : The  $j$ th eigenvalue of (2.1).

$\lambda_{j,h}$ : The  $j$ th eigenvalue of (2.4) computed by the CR finite element.

$\lambda_{j,h}^c$ : The approximation obtained by correcting  $\lambda_{j,h}$ .

$t(s)$ : The CPU time to compute eigenvalues on the finest meshes.

**4.1. Numerical results on  $\Omega \subset \mathbb{R}^2$ .** In this subsection, we present two numerical examples. The first is  $\alpha = \beta = 1$ , the second is  $\alpha = 10 \sin^2(x_1 + x_2) + \frac{1}{6}$  and

$\beta = e^{(x_1-1/2)(x_2-1/2)}$ . We compute on the unit square  $(0, 1)^2$  ( $h_0 = \sqrt{2}$ ), on the L-shaped domain  $(-1, 1)^2 \setminus ([0, 1) \times (-1, 0])$  ( $h_0 = 2\sqrt{2}$ ) and on the regular hexagon with the side length of 1 ( $h_0 = 2$ ); for convenience, we refer to the domains as **S**, **L** and **H**, respectively.




Figure 1. The error curves of the first eigenvalues on the unit square:  $\alpha = \beta = 1$  (left) and  $\alpha = 10 \sin^2(x_1 + x_2) + \frac{1}{6}$ ,  $\beta = e^{(x_1-1/2)(x_2-1/2)}$  (right). Vertical axis: The relative error of eigenvalue. Horizontal axis: The diameter of meshes.




Figure 2. The error curves of the first eigenvalues on the L-shaped domain:  $\alpha = \beta = 1$  (left) and  $\alpha = 10 \sin^2(x_1 + x_2) + \frac{1}{6}$ ,  $\beta = e^{(x_1-1/2)(x_2-1/2)}$  (right). Vertical axis: The relative error of eigenvalue. Horizontal axis: The diameter of meshes.

In order to obtain asymptotic lower bounds for the problem (1.1), we use (3.5) to correct  $\lambda_{1,h}$ . The error curves are depicted in Figures 1–3. New approximate eigenvalues  $\lambda_{1,h}^c$  are listed in Tables 1 and 2. From Figures 1–3 we can see that on each domain, the error curves of  $\lambda_{1,h}^c$  and  $\lambda_{1,h}$  are almost parallel to the line with slope 2, which indicates that  $\lambda_{1,h}^c$  and  $\lambda_{1,h}$  have the same and optimal convergence

order  $\mathcal{O}(h^2)$ . This result coincides with the conclusion of Theorem 3.2. In addition, we can assume that the eigenfunctions corresponding to  $\lambda_1$  are smooth. From Tables 1 and 2, on the one hand, we see that  $\lambda_{1,h}$  converge to  $\lambda_1$  from above and the corrected eigenvalues  $\lambda_{1,h}^c$  converge to  $\lambda_1$  from below, which indicates that the correction (3.5) can provide lower bounds for eigenvalues even though eigenfunctions are smooth. This result coincides with the conclusion of Theorem 3.1. On the other hand, on each domain, the CPU time to compute  $\lambda_{1,h}^c$  is almost the same as that of  $\lambda_{1,h}$ , which tells us that the correction takes very little time.




Figure 3. The error curves of the first eigenvalues on the regular hexagon:  $\alpha = \beta = 1$  (left) and  $\alpha = 10 \sin^2(x_1 + x_2) + \frac{1}{6}$ ,  $\beta = e^{(x_1-1/2)(x_2-1/2)}$  (right). Vertical axis: The relative error of eigenvalue. Horizontal axis: The diameter of meshes.

Domain	S		L		H	
$h$	$\lambda_{1,h}$	$\lambda_{1,h}^c$	$\lambda_{1,h}$	$\lambda_{1,h}^c$	$\lambda_{1,h}$	$\lambda_{1,h}^c$
$\frac{h_0}{32}$	0.24008533	0.24006902	0.34143156	0.34134357	0.39334226	0.39329159
$\frac{h_0}{64}$	0.24008065	0.24007657	0.34141986	0.34139787	0.39332055	0.39330788
$\frac{h_0}{128}$	0.24007948	0.24007846	0.34141699	0.34141149	0.39331513	0.39331196
$\frac{h_0}{256}$	0.24007918	0.24007893	0.34141628	0.34141490	0.39331377	0.39331298
$\frac{h_0}{512}$	0.24007911	0.24007905	0.34141610	0.34141576	0.39331344	0.39331324
$t(s)$	31.10	31.20	22.74	22.81	25.34	25.41
Trend	↘	↗	↘	↗	↘	↗

Table 1. The uncorrected eigenvalues and the corrected eigenvalues on  $\Omega \subset \mathbb{R}^2$ :  $\delta = \frac{100}{99}$ ,  $\alpha = \beta = 1$ .

Domain	<b>S</b>		<b>L</b>		<b>H</b>	
$h$	$\lambda_{1,h}$	$\lambda_{1,h}^c$	$\lambda_{1,h}$	$\lambda_{1,h}^c$	$\lambda_{1,h}$	$\lambda_{1,h}^c$
$\frac{h_0}{32}$	0.24696	0.23963	0.53724	0.27358	0.56181	0.45333
$\frac{h_0}{64}$	0.24645	0.24441	0.51661	0.39617	0.55267	0.51468
$\frac{h_0}{128}$	0.24623	0.24571	0.50301	0.46469	0.54766	0.53740
$\frac{h_0}{256}$	0.24616	0.24603	0.49700	0.48710	0.54600	0.54341
$\frac{h_0}{512}$	0.24614	0.24611	0.49528	0.49280	0.54556	0.54491
$t(s)$	37.71	40.05	28.06	29.74	28.39	30.07
Trend	$\searrow$	$\nearrow$	$\searrow$	$\nearrow$	$\searrow$	$\nearrow$

Table 2. The uncorrected eigenvalues and the corrected eigenvalues on  $\Omega \subset \mathbb{R}^2$ :  $\delta = \frac{100}{99}$ ,  $\alpha = 10 \sin^2(x_1 + x_2) + \frac{1}{6}$ ,  $\beta = e^{(x_1 - 1/2)(x_2 - 1/2)}$ .

Domain	<b>C</b>			<b>F</b>			
$h$	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{5,h}$	$h$	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$
0.6124	0.162344	1.11356	1.56489	0.8660	0.268747	0.54947	0.72763
0.3062	0.162226	1.14537	1.65619	0.4330	0.268359	0.56641	0.73377
0.1531	0.162196	1.15272	1.68222	0.2165	0.268268	0.57235	0.73615
0.0765	0.162189	1.15448	1.68924	0.1083	0.268247	0.57441	0.73687
Trend	$\searrow$	$\nearrow$	$\nearrow$	-	$\searrow$	$\nearrow$	$\nearrow$

Table 3. The CR finite element eigenvalue approximations on  $\Omega \subset \mathbb{R}^3$ :  $\delta = \frac{100}{99}$ .

Domain	<b>C</b>			<b>F</b>		
$h$	$\lambda_{1,h}$	$\lambda_{1,h}^c$	$h$	$\lambda_{1,h}$	$\lambda_{1,h}^c$	
0.6124	0.162344	0.156854	0.8660	0.268747	0.244062	
0.3062	0.162226	0.160802	0.4330	0.268359	0.261752	
0.1531	0.162196	0.161837	0.2165	0.268268	0.266587	
0.0765	0.162189	0.162099	0.1083	0.268247	0.267824	
$t(s)$	150.07	150.23	-	223.12	223.26	
Trend	$\searrow$	$\nearrow$	-	$\searrow$	$\nearrow$	

Table 4. The uncorrected eigenvalues and the corrected eigenvalues on  $\Omega \subset \mathbb{R}^3$ :  $\delta = \frac{100}{99}$ .

**4.2. Numerical results on  $\Omega \subset \mathbb{R}^3$ .** In this subsection, we select  $\alpha = \beta = 1$ . We compute in the cube  $(0, 1)^3$  and the Fichera corner domain  $(-1, 1)^3 \setminus (-1, 0]^3$ . For convenience, we denote the domains by **C** and **F**, respectively. The quasi-uniform mesh samples of the cube and the Fichera corner domain are depicted in Figure 4. In the two domains, we compute the first three eigenvalues using the CR finite element



and list the results in Table 3. In the cube,  $\lambda_2$  and  $\lambda_5$  are the eigenvalues with multiplicity of 3. Corrected eigenvalues  $\lambda_{1,h}^c$  are listed in Table 4. Error curves are depicted in Figure 5.




Figure 4. The quasi-uniform mesh samples of the cube (left) and the Fichera corner domain (right).




Figure 5. The error curves of the first eigenvalues in the cube (left) and the Fichera corner domain (right). Vertical axis: The relative error of eigenvalue. Horizontal axis: The diameter of meshes.

From Figure 5, we see that the error curves of  $\lambda_{1,h}^c$  and  $\lambda_{1,h}$  are parallel to the line with slope 2, which indicates that  $\lambda_{1,h}^c$  and  $\lambda_{1,h}$  have the same and optimal convergence order  $\mathcal{O}(h^2)$ . Also we assume that the eigenfunctions corresponding to  $\lambda_1$  are smooth. From Table 3, we see that on each domain,  $\lambda_{1,h}$  converge to  $\lambda_1$  from above. This shows that the CR finite element eigenvalue approximations may not be lower bounds of exact eigenvalues in the case of smooth eigenfunctions. From Table 4, we see that the corrected eigenvalues  $\lambda_{1,h}^c$  converge to  $\lambda_1$ , which indicates that the correction (3.5) provides lower bounds for eigenvalues even though the eigenfunctions are smooth. The numerical results on three dimensional domains coincide with the conclusions of Theorem 3.1 and Theorem 3.2.

Acknowledgements. We cordially thank the editor and the referees for their valuable comments and suggestions which led to the improvement of this paper.

### References

- [1] *A. Alonso, A. Dello Russo*: Spectral approximation of variationally-posed eigenvalue problems by nonconforming methods. *J. Comput. Appl. Math.* *223* (2009), 177–197. [zbl](#) [MR](#) [doi](#)
- [2] *M. G. Armentano, R. G. Durán*: Asymptotic lower bounds for eigenvalues by nonconforming finite element methods. *ETNA, Electron. Trans. Numer. Anal.* *17* (2004), 93–101. [zbl](#) [MR](#)
- [3] *I. Babuška, J. Osborn*: Eigenvalue problems. *Finite Element Methods (Part 1)*. Handbook of Numerical Analysis II. North-Holland, Amsterdam, 1991, pp. 641–787. [zbl](#) [MR](#)
- [4] *D. Boffi*: Finite element approximation of eigenvalue problems. *Acta Numerica* *19* (2010), 1–120. [zbl](#) [MR](#) [doi](#)
- [5] *J. H. Bramble, J. E. Osborn*: Approximation of Steklov eigenvalues of non-selfadjoint second order elliptic operators. *The Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations* (A. K. Aziz, ed.). Academic Press, New York, 1972, pp. 387–408. [zbl](#) [MR](#) [doi](#)
- [6] *S. C. Brenner, L. R. Scott*: *The Mathematical Theory of Finite Element Methods*. Texts in Applied Mathematics 15. Springer, Berlin, 2002. [zbl](#) [MR](#) [doi](#)
- [7] *C. Carstensen, D. Gallistl*: Guaranteed lower eigenvalue bounds for the biharmonic equation. *Numer. Math.* *126* (2014), 33–51. [zbl](#) [MR](#) [doi](#)
- [8] *C. Carstensen, J. Gedicke*: Guaranteed lower bounds for eigenvalues. *Math. Comput.* *83* (2014), 2605–2629. [zbl](#) [MR](#) [doi](#)
- [9] *C. Carstensen, J. Gedicke, D. Rim*: Explicit error estimates for Courant, Crouzeix-Raviart and Raviart-Thomas finite element methods. *J. Comput. Math.* *30* (2012), 337–353. [zbl](#) [MR](#) [doi](#)
- [10] *I. Chavel, E. A. Feldman*: An optimal Poincaré inequality for convex domains of non-negative curvature. *Arch. Ration. Mech. Anal.* *65* (1977), 263–273. [zbl](#) [MR](#) [doi](#)
- [11] *L. Chen*: iFEM: an innovative finite element methods package in MATLAB. Technical Report, University of California, Irvine, 2008. Available at <https://pdfs.semanticscholar.org/b841/653da0c77051e91f411d4363afe3727f5cc5.pdf>.
- [12] *M. Crouzeix, P.-A. Raviart*: Conforming and nonconforming finite element methods for solving the stationary Stokes equations. I. *Rev. Franc. Automat. Inform. Rech. Operat.* *7* (1973), 33–75. [zbl](#) [MR](#) [doi](#)
- [13] *A. Dello Russo, A. E. Alonso*: A posteriori error estimates for nonconforming approximations of Steklov eigenvalue problems. *Comput. Math. Appl.* *62* (2011), 4100–4117. [zbl](#) [MR](#) [doi](#)
- [14] *E. M. Garau, P. Morin*: Convergence and quasi-optimality of adaptive FEM for Steklov eigenvalue problems. *IMA J. Numer. Anal.* *31* (2011), 914–946. [zbl](#) [MR](#) [doi](#)
- [15] *J. Hu, Y. Huang*: Lower bounds for eigenvalues of the Stokes operator. *Adv. Appl. Math. Mech.* *5* (2013), 1–18. [zbl](#) [MR](#) [doi](#)
- [16] *J. Hu, Y. Huang, Q. Lin*: Lower bounds for eigenvalues of elliptic operators: by nonconforming finite element methods. *J. Sci. Comput.* *61* (2014), 196–221. [zbl](#) [MR](#) [doi](#)
- [17] *J. Hu, Y. Huang, R. Ma*: Guaranteed lower bounds for eigenvalues of elliptic operators. *J. Sci. Comput.* *67* (2016), 1181–1197. [zbl](#) [MR](#) [doi](#)
- [18] *Y. Li*: Lower approximation of eigenvalues by the nonconforming finite element method. *Math. Numer. Sin.* *30* (2008), 195–200. (In Chinese.) [zbl](#) [MR](#)
- [19] *Q. Li, Q. Lin, H. Xie*: Nonconforming finite element approximations of the Steklov eigenvalue problem and its lower bound approximations. *Appl. Math., Praha* *58* (2013), 129–151. [zbl](#) [MR](#) [doi](#)

- [20] *Q. Li, X. Liu*: Explicit finite element error estimates for nonhomogeneous Neumann problems. *Appl. Math., Praha* 63 (2018), 367–379. [zbl](#) [MR](#) [doi](#)
- [21] *Q. Lin, H.-T. Huang, Z.-C. Li*: New expansions of numerical eigenvalues for  $-\Delta u = \lambda \rho u$  by nonconforming elements. *Math. Comput.* 77 (2008), 2061–2084. [zbl](#) [MR](#) [doi](#)
- [22] *Q. Lin, H. Xie*: Recent results on lower bounds of eigenvalue problems by nonconforming finite element methods. *Inverse Probl. Imaging* 7 (2013), 795–811. [zbl](#) [MR](#) [doi](#)
- [23] *Q. Lin, H. Xie, F. Luo, Y. Li, Y. Yang*: Stokes eigenvalue approximations from below with nonconforming mixed finite element methods. *Math. Pract. Theory* 40 (2010), 157–168. [MR](#)
- [24] *X. Liu*: A framework of verified eigenvalue bounds for self-adjoint differential operators. *Appl. Math. Comput.* 267 (2015), 341–355. [zbl](#) [MR](#) [doi](#)
- [25] *F. Luo, Q. Lin, H. Xie*: Computing the lower and upper bounds of Laplace eigenvalue problem: by combining conforming and nonconforming finite element methods. *Sci. China, Math.* 55 (2012), 1069–1082. [zbl](#) [MR](#) [doi](#)
- [26] *J. T. Oden, J. N. Reddy*: An Introduction to the Mathematical Theory of Finite Elements. Pure and Applied Mathematics. Wiley-Interscience, New York, 1976. [zbl](#) [MR](#)
- [27] *G. Savaré*: Regularity results for elliptic equations in Lipschitz domains. *J. Funct. Anal.* 152 (1998), 176–201. [zbl](#) [MR](#) [doi](#)
- [28] *I. Šebestová, T. Vejchodský*: Two-sided bounds for eigenvalues of differential operators with applications to Friedrichs, Poincaré, trace, and similar constants. *SIAM J. Numer. Anal.* 52 (2014), 308–329. [zbl](#) [MR](#) [doi](#)
- [29] *M. Xie, H. Xie, X. Liu*: Explicit lower bounds for Stokes eigenvalue problems by using nonconforming finite elements. *Japan J. Ind. Appl. Math.* 35 (2018), 335–354. [zbl](#) [MR](#) [doi](#)
- [30] *Y. Yang, J. Han, H. Bi, Y. Yu*: The lower/upper bound property of the Crouzeix-Raviart element eigenvalues on adaptive meshes. *J. Sci. Comput.* 62 (2015), 284–299. [zbl](#) [MR](#) [doi](#)
- [31] *Y. Yang, Q. Li, S. Li*: Nonconforming finite element approximations of the Steklov eigenvalue problem. *Appl. Numer. Math.* 59 (2009), 2388–2401. [zbl](#) [MR](#) [doi](#)
- [32] *Y. Yang, Q. Lin, H. Bi, Q. Li*: Eigenvalue approximations from below using Morley elements. *Adv. Comput. Math.* 36 (2012), 443–450. [zbl](#) [MR](#) [doi](#)
- [33] *Y. Yang, Y. Zhang, H. Bi*: A type of adaptive  $C^0$  non-conforming finite element method for the Helmholtz transmission eigenvalue problem. *Comput. Methods Appl. Mech. Eng.* 360 (2020), Article ID 112697, 20 pages. [zbl](#) [MR](#) [doi](#)
- [34] *Y. Yang, Z. Zhang, F. Lin*: Eigenvalue approximation from below using non-conforming finite elements. *Sci. China, Math.* 53 (2010), 137–150. [zbl](#) [MR](#) [doi](#)
- [35] *C. You, H. Xie, X. Liu*: Guaranteed eigenvalue bounds for the Steklov eigenvalue problem. *SIAM J. Numer. Anal.* 57 (2019), 1395–1410. [zbl](#) [MR](#) [doi](#)
- [36] *Z. Zhang, Y. Yang, Z. Chen*: Eigenvalue approximation from below by Wilson’s element. *Math. Numer. Sin.* 29 (2007), 319–321. (In Chinese.) [zbl](#) [MR](#)

*Authors’ addresses:* Yu Zhang, School of Mathematical Sciences, Guizhou Normal University, No. 116 Baoshan Road (N), Guiyang 550001, China; School of Mathematics & Statistics, Guizhou University of Finance and Economics, Huayan Rd, Huaxi District, Guiyang 550025, China, e-mail: zhang\_hello\_hi@126.com; Hai Bi, Yidu Yang (corresponding author), School of Mathematical Sciences, Guizhou Normal University, No. 116 Baoshan Road (N), Guiyang 550001, China, e-mail: bihaimath@gznu.edu.cn, ydyang@gznu.edu.cn.